



RedspiderStop

Reg. No. / Nr. L6747 Act No. / Wet Nr. 36 of / van 1947

GROUP 12D INSECTICIDE

An emulsifiable concentrate for the control of red mite in citrus and red spider mite on apples, pears, peaches, ornamentals, flowers, and cotton, as well as purple, yellow, and flat mite (*Brevipalpus* mite) on tea.

‘n Emulgeerbare konsentraat vir die beheer van rooimyt op sitrus, rooispinmyt op appels, pere, perskes, sierplante, blomme, en katoen, asook pers-, geel-, en platmyt (*Brevipalpus* myt) op tee.

Active Ingredients / Aktiewe Bestanddele:

Tetradifon / Tetradifon 81 g/ℓ



Registered by / Geregistreer deur:

Small Pack Solutions (Pty.) Ltd.

Reg. No. 1997/17900/07

18 Aschenberg Street | Chamdor | Krugersdorp

+27 11 769 1300

Griffon Poison Information Centre: +27 82 446 8946

BATCH NO.:

LOT NR.:

DATE OF MANUFACTURE: See printed information on container

VERVAARDIGINGSDATUM:

EXPIRY DATE:

VERVALDATUM:

UN NO.: 2995

VN NR.:

Content
Inhoud



IMPORTANT / BELANGRIK:

Read label before use /

Lees etiket voor gebruik



**CAUTION
VERSIGTIG**



WARNINGS:

- Allow the following number of days between application and harvest:

Peaches: 28 days

Tea: 21 days

Apples: 14 days

Pears: 14 days

- Store in a cool place away from food and feed.
- May cause eye irritation.
- Keep out of reach of children, uninformed persons, and animals.
- Harmful if swallowed, by contact, or inhalation.
- Toxic to fish.
- **FLAMMABLE** – Do not store or use near open flame.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions. The action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on label, the occurrence of resistance of the pest against the remedy concerned as well as the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or lack of performance of the remedy due to the failure of the user to follow the instructions or to the occurrence of conditions which could not have been foreseen in terms of registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Avoid inhalation of spray mist and contact with skin and eyes.
- Wash contaminated clothing after the day's work.
- Wash with soap and water after accidental skin contact.
- Do not eat, drink, or smoke whilst mixing or applying or before washing hands and face.
- Avoid drift of spray mist onto other crops, grazing, rivers, dams, and areas not under treatment.
- Clean applicator after use and dispose of wash water where it will not contaminate crops, rivers, and dams.
- Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank. Destroy empty container by perforation and flattening and do not re-use for any other purpose.
- Prevent contamination of food, feed, eating utensils and drinking water.

DIRECTIONS FOR USE: Use only as directed.

REDSPIDERSTOP is mainly an ovicide with long residual action and should be applied as a full cover spray as soon as mite infestation is noticed, thus killing eggs and larvae present on trees and plants, and avoiding new deposits of fertile eggs by its action on the adult female mites. It is necessary to apply **REDSPIDERSTOP** early in the season prior to peak mite infestation.

REDSPIDERSTOP is non-toxic to bees and other useful insects and does not kill natural enemies of mites. Spraying may be carried out during blossoming time.

Mixing instructions:

Mix the recommended amount of **REDSPIDERSTOP** into a tank containing the required amount of water whilst stirring. Agitation of the spray liquid during spraying is advisable.

CROP	PEST	DILUTION	REMARKS
APPLES PEARS PEACHES ORNAMENTALS & FLOWERS	Two-spotted & Red spider mite	200 ml/100 ℓ water	Apply as a full cover spray. Repeat at 2 – 3 week intervals.
CITRUS	Red mite	200 ml/100 ℓ water	Apply depending on tree sizes at the rate of 800 – 1000 ℓ spray mix per hectare as full cover spray. Repeat at 2 – 3 week intervals.
COTTON	Red spider mite	2 ℓ/ha	Apply in the case of cotton taller than 60 cm in at least 200 ℓ water/ha. Repeat at 2 – 3 week intervals.
TEA	Purple mite Yellow mite Flat mite (<i>Brevipalpus</i> mite)	400 ml/100 ℓ water	Apply as full cover spray. Repeat at 2 – 3 week intervals.