



# Trojan Ant Bait

Reg. No. / Nr. L8999 Act No. / Wet Nr. 36 of / van 1947

**GROUP 20A INSECTICIDE**

Ready-to-use granular bait for the control of Ants (including the Black and Brown House Ants as well as Argentine Ants) outdoors, around domestic premises, buildings, on lawns, turfgrass, and other areas such as parks and golf courses.

‘n Gereed-vir-gebruik lokaas vir die beheer van Miere (insluitende die Swart en Bruin Huismiere asook Argentynse Miere) vir buitenshuis asook binnenshuis, rondom bewoonde persele, geboue, grasperke, sportvelde, en ander areas soos parke en golfbane.

**Active Ingredients / Aktiewe Bestanddele:**

Hydramethylnon / Hidrametiëlnoon ..... 12 g/kg  
(Amidinohidrazone) / (Amidinohidrasoon)



Registered and Distributed by / Geregistree en Versprei deur:  
**AVIMA (Pty.) Ltd.**  
Reg. No. 1961/001744/07  
18 Aschenberg Street | Chamdor | Krugersdorp  
+27 11 769 1300

Griffon Poison Information Centre: +27 82 446 8946

BATCH NO.:  
LOT NR.:

DATE OF MANUFACTURE: See printed information on container  
VERVAARDIGINGSDATUM:

EXPIRY DATE:  
VERVALDATUM:

UN NO.: 3077  
VN NR.:

Content  
Inhoud

**g**



IMPORTANT / BELANGRIK:  
Read label before use /  
Lees etiket voor gebruik



**CAUTION  
VERSIGTIG**



**WARNINGS:**

- Avoid all contact by ingestion.
- Avoid skin contact.
- Toxic to fish.
- Keep out of reach of children, animals, and uninformed persons.
- Store in a cool, dry place away from food and feedstuffs.
- Do not allow to freeze and keep dry.
- Do not store in direct sunlight.
- Keep container closed when not in use.

**PRECAUTIONS:**

- Do not eat, drink, or smoke while applying the bait.
- Apply only in positions inaccessible to children and animals.
- Wash hands and exposed skin with soap and water, before meals and after work.
- Do not apply the granules to surfaces on which food or feed is stored, prepared, or eaten.
- Protect food preparing equipment and eating utensils from contamination during application.
- Avoid contamination with other insecticides.
- Destroy empty container and NEVER re-use for any other purpose.

**DIRECTIONS FOR USE:** Use only as indicated.

- **TROJAN ANT BAIT** granules exhibit a delayed action and this allows time for transfer of the bait within the colony before the ants die. It is readily transferred from one ant to another through normal food exchange between the ants within the colony. **TROJAN ANT BAIT** will be carried back to the nests by foraging workers before it takes effect.
  - When applied as directed, the bait will control many common structure-invading ants before they enter a building.
  - Reduced activity and dead ants around the entrances of their nests should be noted around 5 – 7 days after baiting.
- NOTE:** Dead ants will not normally be seen where ants are foraging.  
Do not apply any other insecticides on top or in close vicinity to the bait as it may reduce the effectiveness.

**Scatter & Ant nest treatment:**

- Sprinkle the granules lightly on lawns and other areas.
- Sprinkle approximately 30 g bait around active ant nests.
- Do not apply bait directly in or on the nest. The nest should be left undisturbed, as ants will become agitated and may not feed on the bait.

**Perimeter treatment:**

- Lightly sprinkle the granules in a band of 0,5 m wide around the perimeter of buildings where ants are active.
- Treat ant trails, cracks, and crevices where ants may enter structures.
- Acceptance of the bait by ants is dependent on their food preferences which may change depending on the requirements of the nest.
- Ants may not be seen to take the bait immediately as they may already be foraging at other food or water sites.