



# Bait Blocks

Reg. No. / Nr. L9291 Act No. / Wet Nr. 36 of / van 1947

A highly active anticoagulant grain bait for the control of Norway Rat, Roof Rat, the House Mouse and Gerbilles.

'n Hoogs aktiewe antibloedstollingslokaas vir die beheer van die Noorweegse Rot, Dak Rot, die Huismuis en Springhaasrotte.

**Active Ingredients / Aktiewe Bestanddele:**

Brodifacoum (Anticoagulant) /  
Brodifakum (Bloedstolweermiddel)..... 0,05 g/kg

**IMPORTANT / BELANGRIK:** Read label before use / Lees etiket voor gebruik

For BATCH NO. / EXPIRY DATE /  
DATE OF MANUFACTURE:  
See side of container

Vir LOT NR. / VERVALIDATUM /  
VERVAARDIGINGSDATA:  
Sien kant van houer

UN NO. / VN NR: 3026

Content / Inhoud

Registered by /Geregistreeer deur:  
Innovative Pest Management (Pty.) Ltd.

Reg. No. 2001/00587/07  
18 Aschenberg Street | Chamdor | Krugersdorp  
+27 11 769 1300

Griffon Poison Information Centre:  
+27 82 446 8946



**HARMFUL  
SKADELIK**



**WARNINGS:**

- Handle with care.
- Prevent access to bait by children and domestic animals.
- Store under lock and key away from food and feedstuffs.
- Poisonous if swallowed and upon prolonged contact with skin.
- Toxic to fish and wildlife.
- In case of poisoning, call a doctor and make this label available to him.

**PRECAUTIONS:**

- Avoid contamination of any food, food utensils, feeds and drinking water.
- Set bait stations where these will be inaccessible to children and domestic animals.
- Wash hands and exposed skin after handling.
- Do not eat, drink or smoke while handling the bait.
- Destroy empty container and do not re-use it for any other purpose.
- Bury dead rodents and any unconsumed bait at the end of treatment.

**SYMPTOMS OF HUMAN POISONING:**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

**FIRST AID TREATMENT:**

- If swallowed, induce vomiting by tickling the back of the throat. Repeat until vomit is clear.
- Take the patient immediately to the nearest physician.

**NOTE TO PHYSICIAN:**

Intravenous Vitamin K1, 10 – 20 mg depending on prothrombin time. In severe cases small transfusions of fresh blood.

**DIRECTIONS FOR USE: Use only as directed.**

- For use in the home, on the farm, and industrial premises.

**RODEX GRAIN BAIT** can kill rats and mice after a single feed.

- Death of rodents will take 4 – 12 days.
- Effective against Warfarin resistant rats and mice.

**Norway and Roof rats:**

- Place **RODEX GRAIN BAIT** in a covered bait station to prevent access by children and domestic animals.
- Place 50 –75 g of **RODEX GRAIN BAIT**. per bait station along walls and active rat runs, in rat burrows and in sheltered areas where rats are known to live or feed.
- Bait points should be inspected and replenished every 7 days until feeding ceases.

**Mice:**

- Place 15 – 20 g **RODEX GRAIN BAIT** per bait station every 2 – 4 m along walls or runways, in nests and in sheltered areas where mice are known to live and feed.
- The single feeding action of **RODEX GRAIN BAIT** is particularly effective against mice with their small appetites and sporadic feeding behaviour.

**Gerbilles:**

- Place 50 – 75 g **RODEX GRAIN BAIT** in protected bait stations, 10 m apart or 15 – 20 g placed into active holes in the colony.