



Liquid Bait

Reg. No. / Nr. L9289 Act No. / Wet Nr. 36 of / van 1947

A highly active anticoagulant liquid bait for the control of the Norway Rat (*Rattus norvegicus*), Roof Rat (*Rattus rattus*) and the House Mouse (*Mus musculus*).

'n Hoogs aktiewe vloeistofkonsentraat bloedstolweermiddel-lokaas vir die beheer van die Noorweegse Rot (*Rattus norvegicus*), Dak Rot (*Rattus rattus*) en die Huismuis (*Mus musculus*).

Active Ingredients / Aktiewe Bestanddele:

Brodifacoum (Anticoagulant) /
Brodifakum (Bloedstolweermiddel)..... 0,75 g/ℓ

IMPORTANT / BELANGRIK: Read label before use / Lees etiket voor gebruik

For BATCH NO. / EXPIRY DATE /
DATE OF MANUFACTURE:
See side of container

Vir LOT NR. / VERVALIDATUM /
VERVAARDIGINGSDATA:
Sien kant van houer

UN NO. / VN NR: 3026

Content / Inhoud

Registered by /Geregistreeer deur:
Innovative Pest Management (Pty.) Ltd.
Reg. No. 2001/00587/07
18 Aschenberg Street | Chamdor | Krugersdorp
+27 11 769 1300

Griffon Poison Information Centre:
+27 82 446 8946



VERY TOXIC
UITERS GIFTIG



WARNINGS:

- Handle with extreme care.
- Prevent access to bait by children and domestic animals.
- Store under lock and key away from food and feedstuffs.
- Poisonous if swallowed and upon prolonged contact with skin.
- Toxic to fish and wildlife.
- In case of poisoning, call a doctor and make this label available to him.

PRECAUTIONS:

- Avoid all contact by mouth.
- Avoid contamination of any edible commodity, feeds, food utensils and drinking water.
- Set bait stations where these will be inaccessible to children and domestic animals.
- Wear rubber gloves and wash hands after handling.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the tank before destroying the container. Destroy empty container and do not re-use it for any other purpose.
- Do not eat, drink or smoke while handling the concentrate.
- Bury remains at end of treatment and burn or bury rodent bodies.

SYMPTOMS OF HUMAN POISONING:

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

FIRST AID TREATMENT:

- If swallowed, induce vomiting by tickling the back part of the throat. Repeat until vomit is clear and free from smell of poison.
- Take the patient IMMEDIATELY to nearest physician.

NOTE TO PHYSICIAN:

Intravenous Vitamin K1, 50 – 150 mg on first day, in divided doses, depending on prothrombin time. Keep under medical supervision until prothrombin times return to normal. In severe cases small transfusions of fresh blood.

DIRECTIONS FOR USE: Use only as directed.

- For use in the home, on the farm, and industrial premises.
- Effective against Warfarin resistant rats and mice

1. INITIAL BAITING FOR HEAVILY INFECTED AREAS:

- Mix 1 part **RODEX LIQUID BAIT** with 15 parts clean water (65 ml to 1 litre water), place mixed liquid poison into liquid bait dispensers and place in areas of infestation, out of reach of children and animals.
- All bait stations are to be inspected daily for the first week and weekly thereafter.
- Permanent baiting should be continued in heavily infested areas using not only **RODEX LIQUID BAIT** but in combination with **RODEX RAT AND MOUSE PELLETS** and **RODEX GRAIN BAIT**.

2. PERMANENT BAITING:

- Mix 1 part **RODEX LIQUID BAIT** with 50 parts clean water (20 ml to 1 litre).
- Place liquid poison into liquid bait dispensers and place in areas of infestation, out of reach of children and animals.
- All bait stations are to be inspected daily for the first week and weekly thereafter.
- Death normally occurs in 4 - 12 days.