



TM



RODENTICIDE

Professional

FOR USE BY REGISTERED  
PEST CONTROL  
OPERATORS ONLY

# Liquid Concentrate

Reg. No. / Nr. L9290 Act No. / Wet Nr. 36 of / van 1947

A highly active anticoagulant liquid bait concentrate for the control of the Norway Rat (*Rattus norvegicus*), Roof Rat (*Rattus rattus*) and the House Mouse (*Mus musculus*).

'n Hoogs aktiewe vloeistofkonsentraat bloedstolweermiddel-lokaas vir die beheer van die Noorweegse Rot (*Rattus norvegicus*), Dak Rot (*Rattus rattus*) en die Huismuis (*Mus musculus*).

**Active Ingredients / Aktiewe Bestanddele:**

Brodifacoum (Anticoagulant) /  
Brodifakum (Bloedstolweermiddel)..... 2,5 g/ℓ

**IMPORTANT / BELANGRIK:** Read label before use / Lees etiket voor gebruik

For BATCH NO. / EXPIRY DATE /  
DATE OF MANUFACTURE:  
See side of container

Vir LOT NR. / VERVALIDATUM /  
VERVAARDIGINGSDATA:  
Sien kant van houer

UN NO. / VN NR: 3026

Content / Inhoud

Registered by /Geregistreer deur:  
Innovative Pest Management (Pty.) Ltd.  
Reg. No. 2001/00587/07  
18 Aschenberg Street | Chamdor | Krugersdorp  
+27 11 769 1300

Griffon Poison Information Centre:  
+27 82 446 8946



DEVELOPED BY  
**IPM**



**VERY TOXIC  
UITERS GIFTIG**



**WARNINGS:**

- HANDLE WITH EXTREME CARE.
- Prevent access to bait by children, domestic animals and uninformed persons.
- Store under lock and key away from food and feedstuffs.
- Poisonous if swallowed and upon prolonged contact with skin.
- Toxic to fish and wildlife.
- In case of poisoning, call a doctor and make this label available to him.

**PRECAUTIONS:**

- Avoid all contact by mouth.
- Avoid contamination of any edible commodity, feeds, food utensils and drinking water.
- Set bait stations where these will be inaccessible to children and domestic animals.
- Wear rubber gloves and wash hands after handling.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the tank before destroying the container. Destroy empty container and do not re-use it for any other purpose.
- Do not eat, drink or smoke while handling the concentrate.
- Bury remains at end of treatment and burn or bury rodent bodies.

**SYMPTOMS OF HUMAN POISONING:**

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

**FIRST AID TREATMENT:**

- If swallowed, induce vomiting by tickling the back part of the throat. Repeat until vomit is clear.
- Take the patient IMMEDIATELY to nearest physician.

**NOTE TO PHYSICIAN:**

Intravenous Vitamin K1, 50 – 150 mg on first day, in divided doses, depending on prothrombin time. Keep under medical supervision until prothrombin times return to normal. In severe cases small transfusions of fresh blood.

**DIRECTIONS FOR USE: Use only as directed.**

- For use in the home, on the farm, and industrial premises.
- Effective against Warfarin resistant rats and mice

## 1. LIQUID BAIT:

- Mix 1 part **RODEX LIQUID CONCENTRATE** with 160 parts clean water (5 ml to 800 ml water), place mixed liquid poison into liquid bait dispensers and place in areas of infestation, out of reach of children and animals.
- All bait stations are to be inspected daily for the first week and weekly thereafter.
- Permanent baiting should be continued in heavily infested areas using not only **RODEX LIQUID CONCENTRATE** but combining with cereal type bait. This can be obtained by the following method below, or by using **RODEX RAT AND MOUSE PELLETS** or **RODEX GRAIN BAIT**.

## 2. MANUFACTURE OF SOLID BAITS:

- Using cereal type carriers such as crushed wheat, crushed maize, maize meal and other suitable dry bait carriers.
- Mix 8 ml **RODEX LIQUID CONCENTRATE** with 12 ml clean water, then add to 1 kg of bait carrier.
- Mix thoroughly.
- For Rat control, Place 30 - 50 g of bait into each bait station for rat control. Bait stations should be sited 5 m apart.
- For Mouse control, each bait station should contain 10 - 20 g of bait and sited 2 m apart.
- Place baits along walls, in rodent burrows and areas of infestation.
- Weekly inspections of bait stations is very important to achieve effective control.
- Death normally occurs in 4 - 12 days.