



# CarbaKil DP

Reg. No. L6746 & G3703 Act No. 36 of 1947

**GROUP 1A INSECTICIDE**

A residual contact and stomach poison for the control of insects on various crops as listed. Also contact remedy for the control of red mites, tampans, and lice on poultry, and cage birds and in poultry houses. Will also control ticks and fleas on dogs and cats.

'n Nawerkende kontak- en maaggif vir die beheer van insekte op verskillende gewasse soos aangedui. Ook kontakmiddel vir die beheer van rooimyte, tampans en luise op hoenders en hokvoëls en in hoenderhuise. Beheer ook bosluise en vlooië op honde en katte.

**Active ingredient / Aktiewe bestanddeel:**

Carbaryl (Carbamate) / Karbaril (Karbamaat) ... 50 g/kg



Registered by / Geregistreer deur:  
**Small Pack Solutions (Pty.) Ltd.**  
Reg. No. 1997/17900/07  
18 Aschenberg Street | Chamdor | Krugersdorp  
+27 11 769 1300

Griffon Poison Information Centre: +27 82 446 8946

BATCH NO.:  
LOT NR.:  
  
DATE OF MANUFACTURE:  
VERVAARDIGINGSDATUM:  
  
EXPIRY DATE:  
VERVALDATUM:  
  
UN NO.:  
VN NR.: **2757**

**Content  
Inhoud**  
  
**g**

IMPORTANT / BELANGRIK:  
Read label before use /  
Lees etiket voor gebruik





# CarbaKil DP

Reg. No. L6746 & G3703 Act No. 36 of 1947

A residual contact and stomach poison for the control of insects on various crops as listed. Also contact remedy for the control of red mites, tampsans, and lice on poultry, and cage birds and in poultry houses. Will also control ticks and fleas on dogs and cats.

'n Nawerkende kontak- en maaggif vir die beheer van insekte op verskillende gewasse soos aangedui. Ook kontakmiddel vir die beheer van rooimyte, tampsans en luise op hoenders en hokvoëls en in hoenderhuise. Beheer ook bosluise en vlooië op honde en katte.

GROUP	1A	INSECTICIDE
-------	----	-------------

## ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

Carbaryl (Carbamate) / Karbariel (Karbamaat) ..... 50 g/kg

Registered by: **Small Pack Solutions** (Pty) Ltd  
Co. Reg. No.: 1997/17900/07

P O Box 3131, Kenmare, 1745

Tel: +27 11 769 1300

**Griffon Poison Information Centre: +27 82 446 8946**



## WARNINGS:

- Allow the following days between the last application and harvest:
- Tobacco:** ..... 14 days
- Vines:** ..... 14 days
- Cotton:** ..... 7 days
- Lucerne:** ..... 1 day
- Poisonous when absorbed through the skin or swallowed or inhaled.
- Toxic to fish, bees, and wildlife.
- Keep out of reach of children, uninformed persons, and animals.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by conditions such as abnormal soil, climatic and storage conditions; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for the lack of performance of the remedy concerned due to the failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

## PRECAUTIONS:

- Do not inhale dust.
- Wash contaminated clothing daily.
- Wash with soap and water after use.
- Avoid drift of dust onto other crops, grazing, rivers, dams, and areas not under treatment.
- Clean applicator before using with other remedies and dispose of water where it will not contaminate crops, grazing, rivers, and dams.

- TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS:**  
Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank.
- Puncture the triple rinsed container and dispose of via an approved collector or recycler ([www.croplife.co.za](http://www.croplife.co.za)).
- Do not bury, burn, or donate the container to any other parties that may use it as a container for food or beverages.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

## RESISTANCE WARNING:

For resistance management purposes **AVI-CARBO DP** is classified as a Group Code 1A insecticide. Any insect population may contain individuals naturally resistant **AVI-CARBO DP** and other Group Code 1A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly.

**AVI-CARBO DP** or any other Group 1A insecticide may not control these resistant insects. To delay insecticide resistance:

- Avoid exclusive repeated use of insecticides from the same insecticide Group Code.
- Alternate or tank mix with products from different insecticide Group Codes,
- Integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management, contact AVIMA (Pty) Ltd.

**DIRECTIONS FOR USE:** Use only as indicated.

**Red Mites and tampsans in buildings:** Dust all poultry houses thoroughly and pay particular attention to the floor, new litter, perches, nests, nest hay, cracks in wood and walls.

**Lice on birds:** Dust under each wing and around vent.

**Ticks and fleas on dogs & cats:** Sprinkle powder lightly on the hair and rub in well. In the case of cats, the powder must be brushed out afterwards. Repeat weekly or as often as necessary.

## Directions for application:

CROP	PEST	DOSAGE kg/ha	REMARKS
COTTON	Red bollworm, Spring bollworm, Stainers, Syagrus beetle, Jassids, Aphids	15 – 20 kg	Full cover application when pest is noticed.
LAWNS	Caterpillars, Crickets	20 kg (20 g/10 m <sup>2</sup> )	Apply when damage is noticed. A damp sack left on the lawn during the night will reveal infestation of lawn caterpillar when removed in the morning. Light full cover application. Water in.
LUCERN	Caterpillars	15 – 20 kg	Full cover application when pest is noticed.
TOBACCO	American Bollworm, Slug	20 kg	Apply when pest is noticed. Direct application to flower head in the case of American bollworm.
VINES	Mealy bug	10 – 30 kg	Full cover application only to prevent build-up of mealy bug population.

