



InsectiDust

Reg. No. L6786 Act No. 36 of 1947

GROUP 1B | 3A INSECTICIDE

A Dusting Powder Insecticide for use in Agriculture and Public Health with stomach and contact poison action for the control of pests as listed on cotton, groundnuts, sorghum, maize, ornamentals, sub-tropical fruits, potatoes, deciduous fruit, grazing, wheat, lucerne, cruciferae, mushrooms, onions, tobacco, beans. Also for the control of flies, fleas and bedbugs.

'n Stuifpoeier Insekmiddel vir gebruik in die Landbou en Openbare Gesondheid met maag- en kontakgif vir die beheer van plae soos aangedui op katoen, grondbootjies, graansorghum, mielies, sierstruik, sub-tropiese vrugte, aartappels, sagte vrugte, weidings, koring, lusern, koolgewasse, sampioene, uie, tabak, boontjies. Ook vir die beheer van vlieë, vlooië en weeluis.

Active ingredients / Aktiewe bestanddele:

Malathion (Organophosphate) / Malation (Organofosfaat) ... 50 g/kg
Pyrethrum / Piretriene 0,4 g/kg



Registered by / Geregistreer deur:
Small Pack Solutions (Pty.) Ltd.
Reg. No. 1997/17900/07
18 Aschenberg Street | Chamdor | Krugersdorp
+27 11 769 1300

Griffon Poison Information Centre: +27 82 446 8946

BATCH NO.:

LOT NR.:

DATE OF MANUFACTURE:

VERVAARDIGINGSDATUM:

EXPIRY DATE:

VERVALDATUM:

UN NO.:

VN NR.: 2783

Content
Inhoud

g

IMPORTANT / BELANGRIK:
Read label before use /
Lees etiket voor gebruik



**CAUTION
VERSIGTIG**



InsectiDust

Reg. No. L6786 Act No. 36 of 1947

A Dusting Powder Insecticide for use in Agriculture and Public Health with stomach and contact poison action for the control of pests as listed on cotton, groundnuts, sorghum, maize, ornamentals, sub-tropical fruits, potatoes, deciduous fruit, grazing, wheat, lucerne, cruciferae, mushrooms, onions, tobacco, beans. Also, for the control of flies, fleas and bedbugs.

'n Stuifpoeier Insekmiddel vir gebruik in die Landbou en Openbare Gesondheid met maag- en kontakgif vir die beheer van plae soos aangedui op katoen, grondbootjies, graansorghum, mielies, sierstruik, subtropiese vrugte, aartappels, sagte vrugte, weidings, koring, lusern, koolgewasse, sampioene, uie, tabak, boontjies. Ook vir die beheer van vlieë, vlooië en weeluis.

GROUP 1B | 3A INSECTICIDE

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL:

Malathion (Organophosphate) / Malation (Organofosfaat) ... 50 g/kg
Pyrethrum / Piretriene 0,4 g/kg

Registered by: **Small Pack Solutions** (Pty) Ltd
Co. Reg. No.: 1997/17900/07

P O Box 3131, Kenmare, 1745

Tel: +27 11 769 1300

Griffon Poison Information Centre: +27 82 446 8946



WARNINGS:

- Do not dust within the following number of days, before harvest
- Other edible crops:** 7 days
- Cucurbits, Tomatoes, Mushrooms:** 1 day
- Do not apply to cucurbits unless plants are dry.
- Do not use where milk is processed or stored.
- Poisonous by contact, swallowing, and inhalation.
- Toxic to fish, bees, and wildlife.
- Store away from food and feedstuffs.
- Keep out of reach of children, uninformed persons, and animals.
- Keep container well closed.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; compatibility with other substances not mentioned on this label and the occurrence of pest resistant to the remedy, as well as the method, time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation and the environment, or harm to man or animal or for the lack of performance of the remedy due to the failure of the user to follow instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Do not inhale dust.
- Wash with soap and water after use.

- Do not eat, drink, or smoke whilst applying, or before washing hands and face or change of clothing.
- Prevent drift onto other crops, grazing, rivers, dams, or any areas not under treatment.
- Prevent contamination of food, feedstuffs, and water.
- TRIPLE RINSE THE EMPTY CONTAINER AS FOLLOWS:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank.
- Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za).
- Do not bury, burn, or donate the container to any other parties that may use it as a container for food or beverages.

RESISTANCE WARNING:

For resistance management purposes **INSECTIDUST** is classified as a Group Code 1B and 3A insecticide. Any insect population may contain individuals naturally resistant to **INSECTIDUST** and other Group Code 1B and 3A insecticide. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly.

INSECTIDUST or any other Group 1B and 3A insecticides may not control these resistant insects. To delay insecticide resistance:

- Avoid exclusive repeated use of insecticides from the same insecticide Group Code.
- Alternate or tank mix with products from different insecticide Group Codes,
- Integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management, contact AVIMA (Pty) Ltd.

DIRECTIONS FOR USE: Use only as indicated.

Apply by hand, knapsack, or power duster.

CROP	PEST	DOSAGE	TIME OF APPLICATION
ALL CROPS	Astylus beetle Lesser armyworm Armyworm Chafer beetle Short-horn grasshopper	15 – 30 kg/ha 15 – 30 kg/ha 15 kg/ha 15 – 25 kg/ha 15 – 30 kg/ha	Apply when pest appears and repeat when necessary.
TOBACCO	Aphids Slugs Whitefly	Seedbed treatment 30 g/10 m ² Field treatment 15 – 25 kg/ha	Full cover application when necessary.
SORGHUM TOMATOES FLOWERS & ORNAMENTALS	Aphids	15 – 30 kg/ha	Full cover application when necessary.
PEACHES PLUMS	Aphids	25 kg/ha	Full cover application when necessary.
GROUNDNUTS MAIZE	Aphids	15 – 30 kg/ha	Full cover application when necessary.
COTTON	Aphids	30 kg/ha	Full cover application when necessary.
BEANS	Aphids CMR beetles Thrips	15 – 25 kg/ha 20 – 25 kg/ha 15 – 30 kg/ha	Full cover application when pest is noticed. Repeat when necessary.
FLOWERS & ORNAMENTALS ONIONS	Thrips	15 – 30 kg/ha 20 – 30 kg/ha	Full application at weekly intervals as soon as pest or damage is noticed.
LUCERNE	Caterpillars	10 – 30 kg/ha	Full cover application when pest is noticed.
MUSHROOMS	Phorid & Sciarid flies (Fungus gnats)	10 – 15 kg/ha	Apply thoroughly soon after picking. Repeat at weekly intervals.
CRUCIFERAE (Cabbage, Cauliflower, Broccoli, Brussel sprouts)	Greater cabbage and Diamondback moth larvae	20 – 30 kg/ha	Full cover application when pest is noticed. Repeat at weekly intervals.

Public Health:

SITE	PEST	DOSAGE	APPLICATION DIRECTIONS
GARBAGE MANURE COMPOST HEAPS	Fly larvae	Apply uniformly until visible.	Full cover application to outside of heaps when fly breeding is noticed. Usually at weekly intervals. Apply again whenever fresh material is added.
BUILDINGS ROOMS HOUSES	Fleas Bedbugs	Dust freely	Do not apply to the body. Apply freely to all places frequented by these pests.